

Recreational Water Quality

Current Status and Opportunities

Richard Whitman

Lake Michigan Ecological Research Station

Great Lakes Science Center



- Substantial inconsistency in monitoring approaches among and within states.
- Inconsistent public notification programs.
- Growing concerns about microbiological contaminants
- Increased pollution pressures on coastal and water-based areas due to population growth and more intense land use.

EPA BEACH Program

- Strengthen beach programs and water quality standards
- Inform public about recreational water quality
- Conduct research to improve science for beach programs

Criteria For Quality Indicator

- Relevance to health effects
- Source characterization
- Timely results
- Data confidence
- Beach specificity
- Economy, ease, effectiveness

EPA/CDC/USGS Pilot Epidemiological Study




Beach Sampling Approach

- Test for E. coli or fecal coliform
- Samples generally taken
 - Morning
 - At shore or knee deep
 - From the surface
- Monthly, weekly, or daily
- Single sample per beach






Sample Sizes Required to Achieve Precision + 30% of the Mean




Based on ten replicate samples taken each day in AM at center transect, 90 cm. Estimates calculated using Elliot's (1977) equation for small sample size.

Day to Day Sampling of *E. coli* at Indiana Dunes National Lakeshore



Little Glen Lake, Sleeping Bear Dunes National Lakeshore (*E. coli* counts in CFU/100 ml)



Correlations among Seagull numbers and E. coli concentrations




63rd St. Beach, Chicago, Summer 2000

	# gulls unlagged, P values	# gulls lagged 1 day, P values
Foreshore sand	0.133	0.000*
Submerged sand	0.972	0.046
45 cm water AM	0.224	0.004*
90 cm water AM	0.037	0.001*
45 cm water PM	0.916	0.167
90 cm water PM	0.432	0.008

Results of Light/Dark Bag Experiment


63rd St. Beach, Chicago September 18, 2000




Treatment A Light Bag A Dark Bag A Ambient Water



E. coli in Sand



E.coli in sand and water of 63rd St. Beach,






Partial R, 45cm water vs Shore Sand = 0.501** Partial R Offshore vs Shore Sand = 0.259*





Linear regression of E. coli concentrations in sand and water


63rd St. Beach, Chicago, Summer, 2000



R sq. = 0.43

Variation in *E. coli* density-2001







* Rainfall event

E. coli Concentration of Lake Water, Pore water and Sand with Rainfall


Lakeview Beach, Indiana Dunes National Lakeshore




E. coli and enterecocci persist in Cladophora mats



Concentration of *E. coli* washed from *Cladophora glomerata*



Stream water and sediments are correlated but not with forest soil




Stream Water-A
Stream Sand-B
Margin Sand-C
Sand @ 1 m from margin-D


Soil @ 4 m from margin-E

Connected Lines Indicate Significant Correlation (Spearman rho, p=0.05, n=15)

E. coli is commonly found in sediments surrounding seeps and springs but not emanating water.



E. coli in Dunes Creek increases with stream order



General Forces

- Currents
- Sediments
- Orientation
- Weather
- Runoff





Local Forces



- Morphology
- Shoreline Exposure
- Sources
 - Direct (e.g. sewage, birds)

Indirect (e.g. sands, rivers, submerged

storage)



Correlations among summer, 2000 E. coli concentrations at three Lake Michigan



beaches




Lake Michigan

West Beach Portage, IN

Future Directions

- Better notification
- More Timely Results
 - Faster tests
 - Chemical and Biological
 - Predictive models
 - Local and regional models
- More interjurisdictional cooperation
- Standardization





Richard Whitman Lake Michigan Ecological Research Station United States Geological Survey Porter, IN

Phone: (219) 926-8336 ext. 424

Fax: (219) 929-5792

Email: richard_whitman@usgs.gov